

## Acute limb ischemia secondary to radiation-induced arteritis: case report

### *Insuficiência arterial aguda secundária a arterite induzida por radiação: relato de caso*

José Emerson dos Santos Souza<sup>1</sup>, Leonardo Pessoa Cavalcante<sup>1</sup>, Marcos Velludo Bernardes<sup>1</sup>,

Marcos Henrique Parisati<sup>1</sup>, Patrícia de Souza Lacerda<sup>1</sup>, Raquel Magalhães Pereira<sup>2</sup>

#### Abstract

Radiation-induced arteritis is a rare but well-known complication of radiotherapy. This report describes the case of a 34-year-old woman with uterine cervical cancer who was diagnosed with left iliofemoral deep vein thrombosis (DVT) 2 years after radiotherapy, and 2 months later, during the treatment of DVT with effective anticoagulation, developed an episode of acute arterial ischemia of the left lower limb secondary to a long subocclusive lesion of the external iliac artery. The patient was treated with angioplasty and stenting of the lesion and recovered uneventfully after the endovascular procedure.

**Keywords:** radiotherapy; constriction, pathologic; angioplasty.

#### Resumo

A arterite induzida por radiação é uma rara mas bem documentada complicação da radioterapia. O presente relato descreve o caso de uma mulher de 34 anos, diagnosticada com neoplasia de colo do útero, a qual, dois anos após sessões de radioterapia desenvolveu trombose venosa profunda (TVP) iliofemoral esquerda; dois meses depois, durante tratamento para TVP com devida anticoagulação, a paciente apresentou quadro de insuficiência arterial aguda do membro inferior esquerdo secundária a uma longa lesão suboclusiva da artéria ilíaca externa. A paciente foi tratada com angioplastia transluminal percutânea e implantação de stent autoexpansível, recuperando-se sem intercorrências após o procedimento endovascular.

**Palavras-chave:** radioterapia; constrição patológica; angioplastia.

<sup>1</sup>Hospital Universitário Francisca Mendes, Vascular and Endovascular Surgery Service, Manaus, AM, Brazil.

<sup>2</sup>Universidade Federal do Amazonas – UFAM, Manaus, AM, Brazil.

Financial support: None.

Conflicts of interest: No conflicts of interest declared concerning the publication of this article.

Submitted on: 11.02.12. Accepted on: 04.13.13.

The study was carried out at Hospital Universitário Francisca Mendes, Manaus (AM), Brazil.





**Figure 2.** Post-dilatation of two self-expandable nitinol stents with a 7-mm × 80-mm non-compliant percutaneous transluminal angioplasty balloon.



**Figure 3.** Completion angiogram shows satisfactory angiographic result.

of patients that undergo pelvic region radiotherapy. Novais et al.<sup>5</sup> found that esophageal radiation-induced stenosis affects 25% to 67% of patients irradiated for the treatment of esophageal primary tumors and 1% to 20% for the treatment of adjacent tumors (breast, lung, thyroid).

Although still rarely seen in oncology practice, arterial radiation-induced stenosis is becoming more

common due to the increased use of radiotherapy and prolonged survival of cancer patients<sup>1,6</sup>. The vessels involved are directly associated with cancer location and the irradiation received.

Arterial occlusive disease pathogenesis begins with direct cellular damage due to radiation or by free radicals produced by its action<sup>7</sup>. Cell damage leads to fibrosis, intimal thickening, elastic middle layer degeneration, adventitia fibrosis and vasa vasorum damage<sup>3,6</sup>. After that, in addition to narrowing, there may be thrombus formation, ulceration and distal embolization<sup>1</sup>.

Although well documented, histological changes are nonspecific and virtually identical to those observed in atherosclerosis<sup>1,2</sup>. Some authors suggest that vasa vasorum injury and the consequent arterial wall ischemia are among the few morphological characteristics that separate radiation-induced lesions from spontaneous atherosclerosis<sup>8</sup>. It has also been suggested that the presence of atherosclerosis risk factors further promotes lesion development after radiotherapy<sup>2,6</sup>. However, the predisposition to atherosclerosis only accelerates vascular injury after irradiation, because the absence of lesions at uninvolved areas indicates that radiotherapy is the primary etiology<sup>1,2,6</sup>.

Therefore, the treatment of cervical cancer was the main cause of arterial injury in our patient, as arteriography showed that the disease was restricted to the left EIA, and all the other pelvic and left lower limb vessels had a normal angiographic appearance. Besides that, age and the absence of other atherosclerosis risk factors confirmed the radiation-induced etiology of the arterial disease. The radiation dose reported in the literature to be associated with iliac and femoral artery stenosis is 39.5 to 80 Gy<sup>9</sup>.

In our patient, acute arterial insufficiency symptoms appeared about 2 years after radiotherapy. In the literature, the time between irradiation and signs and symptoms of damage varies: Piedbois et al.<sup>2</sup> described a range of 1 to 10 years (mean 6 years); Dorresteijn et al.<sup>10</sup>, reported a mean 13 years; Hassen-Khodja et al.<sup>11</sup> found a mean 10 years between cervical irradiation and carotid stenosis treatment.

The clinical signs and symptoms of radiation-induced arterial stenosis are not different from other chronic arterial blockages. Therefore, in the arterial involvement of the lower extremities, varying degrees of arterial insufficiency may occur, from exertion muscular ischemia (intermittent claudication) to rest pain and tissue loss<sup>12</sup>.

Our patient presented with an uncommon clinical picture of acute arterial insufficiency. Although

she had a chronic arterial lesion that was probably asymptomatic, iliofemoral DVT may have led to an imbalance of the arterial blood supply to the limb, previously compensated through collateral arteries. These arteries may explain why the pulses were palpable during physical examination two months before acute ischemia.

The clinical diagnosis of chronic arterial insufficiency is usually confirmed using color Doppler US and physiological tests (such as ABI)<sup>13,14</sup>. Digital subtraction arteriography is usually reserved for surgical planning<sup>1,2,10</sup>. In the case described here, urgent arteriography was performed after the clinical diagnosis of acute arterial occlusion, and the endovascular treatment was administered at the same time.

Although other treatment options have been described (drug therapy, endarterectomy, extra-anatomic bypass)<sup>1,2</sup>, PTA with stent implantation is increasingly becoming the treatment of choice in similar cases because of its low invasiveness and high success rates, as described in the literature<sup>12,14</sup>. PTA associated with stent implantation is extremely effective in the treatment of iliac arterial stenosis of any etiology<sup>14</sup>. Patients usually have a good postoperative recovery, and clinical improvement is evident, as shown in this case and in the literature<sup>12,14</sup>.

## CONCLUSION

This case demonstrated that PTA with implantation of self-expandable stents is a good therapeutic option for the treatment of radiation-induced EIA stenosis with clinical symptoms in patients treated for advanced pelvic cancer.

## REFERENCES

- Baerlocher MO, Rajan DK, Douglas J, Rubin BB. Primary stenting of bilateral radiation-induced external iliac stenoses. *J Vasc Surg.* 2004;40:1028-31. PMID:15557921. <http://dx.doi.org/10.1016/j.jvs.2004.08.031>
- Piedbois P, Becquemin JP, Blanc I, et al. Arterial occlusive disease after radiotherapy: a report of fourteen cases. *Radiother Oncol.* 1990;17:133-40. [http://dx.doi.org/10.1016/0167-8140\(90\)90101-2](http://dx.doi.org/10.1016/0167-8140(90)90101-2)
- Warrington KJ, Cooper Jr LT. Vasculitis and Other Arteriopathies. In: Cronenwett JL, Johnston KW, editors. *Rutherford's Vascular Surgery*. Philadelphia: WB Saunders; 2010. p. 1156-68. <http://dx.doi.org/10.1016/B978-1-4160-5223-4.00076-7>
- Teixeira FV, Pilon B, Marchioni R. Tratamento da retite actínica hemorrágica com o uso de solução de formalina intra-retal: Relato de caso. *Rev Bras Coloproct.* 2002;22(3):184-9.
- Novais P, Lemme E, Equi C, Medeiros C, Lopes C, Vargas C. Estenoses benignas de esôfago: abordagem endoscópica com velas de Savary-Gilliard. *Arq Gastroenterol.* 2008;45(4):290-4. PMID:19148356. <http://dx.doi.org/10.1590/S0004-28032008000400006>
- Farrugia M, Gowda KMS, Cheattle TR, Ashok TP. Radiotherapy-Related Axillary Artery Occlusive Disease: Percutaneous Transluminal Angioplasty and Stenting. Two Case Reports and Review of the Literature. *Cardiovasc Intervent Radiol.* 2006;29:1144-7. PMID:16845557. <http://dx.doi.org/10.1007/s00270-005-0230-x>
- Brown KR, Rzucidlo E. Acute and chronic radiation injury. *J Vasc Surg.* 2011;53(15S):15S-21S. PMID:20843630. <http://dx.doi.org/10.1016/j.jvs.2010.06.175>
- Zhou W, Bush RL, Lin PH, Lumsden AB. Radiation-associated venous stenosis: Endovascular treatment options. *J Vasc Surg.* 2004;40(1):179-82. PMID:15218482. <http://dx.doi.org/10.1016/j.jvs.2004.03.039>
- Himmel PD, Hassett JM. Radiation-induced chronic arterial injury. *Semin Surg Oncol.* 1986;2:225-47. <http://dx.doi.org/10.1002/ssu.2980020405>
- Dorresteijn LDA, Vogels OJM, Leeuw FE, et al. Outcome of carotid artery stenting for radiation-induced stenosis. *Int. J. Radiation Oncology Biol. Phys* 2010;77(5):1386-90. PMID:20116932. <http://dx.doi.org/10.1016/j.ijrobp.2009.06.045>
- Hassen-Khodja R, Sala F, Declémy S, Lagrange JL, Bouillane PJ, Batt M. Surgical Management of Atherosclerotic Carotid Artery Stenosis after Cervical Radiation Therapy. *Ann Vasc Surg.* 2000;14(6):608-11. <http://dx.doi.org/10.1007/s100169910110>
- Pedron C, Ristow A, Cury JM Fº, Martin HS, Peixoto CC, Fonseca LMB. Tratamento endovascular da oclusão das artérias ilíacas. *Radiol Bras.* 2001;34(5):261-5. <http://dx.doi.org/10.1590/S0100-39842001000500004>
- Donas KP, Schwindt A, Pitoulias GA, Schönefeld T, Basner C, Torsello G. Endovascular treatment of internal iliac artery obstructive disease. *J Vasc Surg.* 2009;49(6):1447-51. PMID:19497505. <http://dx.doi.org/10.1016/j.jvs.2009.02.207>
- Sullivan TM, Childs MB, Bacharach JM, Gray BH, Piedmonte ML. Percutaneous transluminal angioplasty and primary stenting of the iliac arteries in 288 patients. *J Vasc Surg.* 1997;25(5):829-39. [http://dx.doi.org/10.1016/S0741-5214\(97\)70212-X](http://dx.doi.org/10.1016/S0741-5214(97)70212-X)

### Correspondence

Leonardo Pessoa Cavalcante  
Hospital Universitário Francisca Mendes – Serviço de Cirurgia Vascular/Endovascular  
Av. Camapuã, 108 - Cidade Nova II  
CEP 69097-720 - Manaus (AM), Brazil  
E-mail: leonardocavalcante@uol.com.br

### Author information

LPC is chief, Endovascular Surgery Service, Hospital Universitário Francisca Mendes.  
MVB is chief, Vascular Surgery Service, Hospital Universitário Francisca Mendes.  
MHP is vascular surgeon, Vascular Surgery Service, Hospital Universitário Francisca Mendes.  
JESS and PSL are resident physicians (Vascular Surgery), Vascular Surgery Service, Hospital Universitário Francisca Mendes.  
RMP is medical student, Universidade Federal do Amazonas (UFAM).

### Author's contributions

Conception and design: JESS, LPC, RMP  
Analysis and interpretation: MHP, MVB  
Data collection: JESS, PSL  
Writing the article: JESS, LPC, RMP  
Critical revision of the article: LPC, MVB, MHP  
Final approval of the article\*: JESS, LPC, MVB, MHP, PSL, RMP  
Statistical analysis: N/A  
Overall responsibility: JESS  
Obtained funding: None.

\*All authors should have read and approved of the final version of the article submitted to *J Vasc Bras.*